arXiv:1205.5199v3 [cs.DM] 1 Dec 2012

Automorphism groups of Cayley graphs generated by connected transposition sets

Ashwin Ganesan*

Abstract

Let S be a set of transpositions that generates the symmetric group S_n, where $n \geq 4$. The transposition graph $T(S)$ is defined to be the graph with vertex set $\{1, \ldots, n\}$ and with vertices i and j being adjacent in $T(S)$ whenever $(i, j) \in S$. We prove that if the girth of the transposition graph $T(S)$ is at least 5, then the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$, where $\mathrm{Aut}(S_n, S)$ is the set of automorphisms of S_n that fixes S. This strengthens a result of Feng on transposition graphs that are trees. We also prove that if the transposition graph $T(S)$ is a 4-cycle, then the set of automorphisms of the Cayley graph $\mathrm{Cay}(S_4, S)$ that fixes a vertex and each of its neighbors is isomorphic to the Klein 4-group and hence is nontrivial. We thus identify the existence of 4-cycles in the transposition graph as being an important factor in causing a potentially larger automorphism group of the Cayley graph.

Index terms — Cayley graphs; transposition sets; automorphism groups; modified bubble-sort graph.

1 Introduction

Let $\Gamma = (V, E)$ be a simple, undirected graph. An automorphism of Γ is a permutation of the vertex set that preserves adjacency; in other words, a permutation $\pi \in \mathrm{Sym}(V)$ is an automorphism of Γ if for every $u, v \in V$, $\{u, v\} \in E$ iff $\{\pi(u), \pi(v)\} \in E$. The set of all automorphisms of Γ forms a permutation group called the automorphism group of Γ, which we denote by $\mathrm{Aut}(\Gamma)$.

Let H be a group and let S be a subset of H. The Cayley digraph of H with respect to S, denoted by $\mathrm{Cay}(H, S)$, is the digraph with vertex set H and with an arc from h to sh whenever $h \in H$ and $s \in S$. When S is closed under inverses, (g, h) is an arc of the Cayley digraph if and only if (h, g) is an arc, and so we can

---

*Department of Mathematics, Amrita School of Engineering, Amrita University, Amritanagar, Coimbatore - 641 112, Tamil Nadu, India. Email: `ashwin.ganesan@gmail.com`, `g_ashwin@cb.amrita.edu`.

identify the two arcs (g,h) and (h,g) with the undirected edge $\{g,h\}$. When $1 \notin S$, the Cayley digraph contains no self-loops. Thus, when $1 \notin S = S^{-1}$, $\mathrm{Cay}(H,S)$ can be considered to be a simple, undirected graph. The Cayley graph $\mathrm{Cay}(H,S)$ is connected if and only if S generates H.

The automorphism group of the Cayley graph $\mathrm{Cay}(H,S)$ contains the right regular representation $R(H)$ as a subgroup, and hence all Cayley graphs are vertex-transitive. Let e denote the identity element of the group H and also the corresponding vertex of $\mathrm{Cay}(H,S)$. Since $R(H)$ is regular, $\mathrm{Aut}(\Gamma) = \mathrm{Aut}(\Gamma)_e R(H)$, where $\mathrm{Aut}(\Gamma)_e$ is the stabilizer of e in $\mathrm{Aut}(\Gamma)$. The set of automorphisms of the group H that fixes S setwise, denoted by $\mathrm{Aut}(H,S) := \{\pi \in \mathrm{Aut}(H) : S^\pi = S\}$, is a subgroup of $\mathrm{Aut}(\Gamma)_e$ (cf. [1]). For any Cayley graph $\Gamma = \mathrm{Cay}(H,S)$, the normalizer $N_{\mathrm{Aut}(\Gamma)}(R(H))$ is equal to the semidirect product $R(H) \rtimes \mathrm{Aut}(H,S)$ (cf. [4], [8]). A Cayley graph $\Gamma := \mathrm{Cay}(H,S)$ is said to be normal if $R(H)$ is a normal subgroup of $\mathrm{Aut}(\Gamma)$, or equivalently, if $\mathrm{Aut}(\Gamma) = R(H) \rtimes \mathrm{Aut}(H,S)$. Thus, normal Cayley graphs are those that have the smallest possible full automorphism group (cf. [8]).

Let S be a set of transpositions in the symmetric group S_n. The transposition graph $T(S)$ is defined to be the graph with vertex set $\{1,2,\ldots,n\}$ and with vertices i and j being adjacent in $T(S)$ whenever $(i,j) \in S$. A set of transpositions of $\{1,\ldots,n\}$ generates S_n if and only if the transposition graph $T(S)$ is connected; thus, a set of transpositions is a minimal generating set for S_n if and only if $T(S)$ is a tree (cf. [5]). In the sequel, we assume that the set of transpositions S generates S_n; equivalently, we assume the graph $T(S)$ is connected.

Let G denote the automorphism group of a graph Γ. For a vertex v of Γ, let G_v denote the stabilizer subgroup of G that fixes the vertex v. Let L_v denote the set of automorphisms of Γ that fixes the vertex v and each of its neighbors. The subgroup L_v is called the stabilizer of the vertex neighborhood, and L_v is a normal subgroup of G_v (cf. [1, Chapter 16]).

While one can often obtain some automorphisms of a graph, it is often difficult to prove that one has obtained the (full) automorphism group. A uniqueness property that has been has been useful in proving that one has obtained the full automorphism group of Cayley graphs $\mathrm{Cay}(S_n,S)$ generated by transpositions is the following: given any two elements $t,k \in S$, there is either a unique 4-cycle or a unique 6-cycle in the Cayley graph $\mathrm{Cay}(S_n,S)$ that contains the three vertices e,t and k. This uniqueness property implies, via a proof by induction on the length of a word in terms of the generating elements S, that any automorphism of the Cayley graph $\mathrm{Cay}(S_n,S)$ that fixes a vertex and each of its neighbors must be trivial, i.e. that $L_e = 1$. This implies that the Cayley graph $\mathrm{Cay}(S_n,S)$ has the smallest possible full automorphism group $R(S_n) \rtimes \mathrm{Aut}(S_n,S)$. In Godsil and Royle [5], the uniqueness of certain 4-cycles in the Cayley graph $\mathrm{Cay}(S_n,S)$ was used to determine the automorphism group of $\mathrm{Cay}(S_n,S)$ for the special case when $T(S)$ is an asymmetric tree. In Zhang and Huang [9], the uniqueness of certain 6-cycles in $\mathrm{Cay}(S_n,S)$ was used to determine the automorphism group of $\mathrm{Cay}(S_n,S)$ for the special case when $T(S)$ is a path graph. These results were generalized in Feng [3] to the case where $T(S)$ is an arbitrary tree.

Let $n \geq 3$. Feng [3] showed that if S is a set of transpositions in S_n, then $\mathrm{Aut}(S_n,S) \cong \mathrm{Aut}(T(S))$. This result holds even if $T(S)$ is not a tree. In the special case when $T(S)$ is a tree, Feng [3] showed that the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$. In this case, the stabilizer L_e of a vertex neighborhood of the Cayley graph $\mathrm{Cay}(S_n, S)$ is trivial.

In the present paper, we generalize and extend the results of Feng on trees mentioned in the previous graph to arbitrary transposition graphs that may contain cycles. The girth of a graph is the length of the shortest cycle in the graph. In particular, trees have infinite girth. We prove that if the girth of the transposition graph is at least 5, then the stabilizer subgroup L_e of the Cayley graph $\mathrm{Cay}(S_n, S)$ is trivial and the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$. On the other hand, if $T(S)$ is a 4-cycle, then L_e is shown to be Klein 4-group and hence is not trivial. These results thus identify 4-cycles in the transposition graph as playing an important role in leading to a potentially larger automorphism group (of the Cayley graph) than otherwise.

The main result of this paper is the following:

Theorem 1. *Let S be a set of transpositions that generates S_n, where $n \geq 4$. If the girth of the transposition graph $T(S)$ is at least 5, then the automorphism group of the Cayley graph $Cay(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$, where $\mathrm{Aut}(S_n, S)$ is the set of automorphisms of S_n that fixes S, and the stabilizer of a vertex neighborhood of $\mathrm{Cay}(S_n, S)$ is trivial. If the transposition graph $T(S)$ is a 4-cycle, then the stabilizer of a vertex neighborhood of $\mathrm{Cay}(S_4, S)$ is isomorphic to the Klein 4-group $\mathbb{Z}_2 \times \mathbb{Z}_2$ and hence is nontrivial.*

Remark 1: A special case of the first part of Theorem 1 is the result of Feng [3] that if the transposition graph $T(S)$ is a tree, then the automorphism group of the Cayley graph $\mathrm{Cay}(S_n, S)$ is the semidirect product $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$. Further special cases of this result are the following: the result of Godsil and Royle [5] that if $T(S)$ is an asymmetric tree, then $\mathrm{Cay}(S_n, S)$ has automorphism group $R(S_n) \cong S_n$; the result of Huang and Zhang [7] that if $T(S)$ is the star $K_{1,n-1}$, then $\mathrm{Cay}(S_n, S)$ has automorphism group isomorphic to $S_n S_{n-1}$; and the result of Zhang and Huang [9] that if $T(S)$ is the path graph then $\mathrm{Cay}(S_n, S)$ has automorphism group isomorphic to $S_n \mathbb{Z}_2$.

Remark 2: There is a flaw in the proof by Zhang and Huang [9] that led to an incorrect result in that paper, and the incorrect result there that was recalled and stated in Feng [3] as well; we now recall these statements. When the transposition graph $T(S)$ is the n-cycle graph, the Cayley graph $\mathrm{Cay}(S_n, S)$ is called the *modified bubble-sort graph* of dimension n. It was claimed in [9] that when $T(S)$ is an n-cycle graph for $n \geq 4$, then 6-cycles satisfying some particular conditions in the Cayley graph $\mathrm{Cay}(S_n, S)$ are unique. This uniqueness implies that any automorphism of the Cayley graph that fixes a vertex and each of its neighbors is trivial. It was stated in [3, p.72] that when $T(S)$ is a 4-cycle, by this uniqueness property, the Cayley graph $\mathrm{Cay}(S_4, S)$ is normal and has full automorphism group $R(S_4) \rtimes \mathrm{Aut}(S_4, S) \cong R(S_4)G_e \cong S_4 D_8$, which contains 192 elements.

In the present paper, it is shown that the 6-cycles satisfying certain conditions in the Cayley graph $\mathrm{Cay}(S_4, S)$ are not unique. In particular, the uniqueness proofs in [9] are incorrect, and computer simulations have confirmed our results as being correct. Thus, the vertex neighborhood stabilizer L_e in this Cayley graph need not be trivial. We determine this stabilizer to be the Klein 4-group $\mathbb{Z}_2 \times \mathbb{Z}_2$. With the help of a computer ([6]) it was confirmed that the full automorphism group of $\mathrm{Cay}(S_4, S)$ has 768 elements and has structure description $(\mathbb{Z}_2 \times \mathbb{Z}_2 \times ((\mathbb{Z}_2^4 \rtimes \mathbb{Z}_3) \rtimes \mathbb{Z}_2)) \rtimes \mathbb{Z}_2$. The full automorphism group of $\mathrm{Cay}(S_4, S)$ obtained in this paper is in a more comprehensible form, i.e. as stated in Theorem 1, it is $G = R(S_4)G_e = R(S_4)(L_e \rtimes D_8) \cong R(S_4)(V_4 \rtimes D_8)$. In this case, $R(S_4)$ is not a normal subgroup of $\mathrm{Aut}(\mathrm{Cay}(S_4, S))$.

2 Proof of Theorem 1

We assume throughout that S is a set of transpositions of $\{1, \ldots, n\}$ that generates S_n. Thus the transposition graph $T(S)$ is connected.

When $n = 3$ and $T(S)$ is the 3-cycle graph, the Cayley graph $\mathrm{Cay}(S_n, S)$ is the complete bipartite graph $K_{3,3}$ (cf. Biggs [1, Chapter 16]). The complement of $K_{3,3}$ is the union of two disjoint 3-cycles. Thus, the automorphism group of $K_{3,3}$ is the automorphism group of a partition of a six element set into two equal sized subsets and hence is equal to the semidirect product $(S_3 \times S_3) \rtimes \mathbb{Z}_2$ (cf. [2, p. 46]).

Thus, the cases where $n \leq 3$ are easily dealt with; we assume in the sequel that $n \geq 4$. We first recall a preliminary result.

Lemma 2. *(Godsil and Royle [5, Lemma 3.10.3]) Let S be a set of transpositions such that the transposition graph $T(S)$ does not contain triangles, and let $t, k \in S$. Then, $tk = kt$ if and only if there is a unique 4-cycle in $\mathrm{Cay}(S_n, S)$ containing e, t and k.*

While the hypothesis in Lemma 2 assumes the transposition graph does not contain triangles, the conclusion in the Lemma holds even if the transposition graph does contain triangles; however, this stronger assertion is not useful in our proof because when the transposition graph $T(S)$ contains triangles, it turns out that the Cayley graph $\mathrm{Cay}(S_n, S)$ does not contain unique 6-cycles satisfying certain conditions. The uniqueness of both 4-cycles and 6-cycles is necessary in order for the proof to go through that L_e is trivial.

We first consider the case where the transposition graph $T(S)$ is an n-cycle graph. In this case, the set of generators S is $\{(1, 2), (2, 3), \ldots, (n-1, n), (n, 1)\}$, and the corresponding Cayley graph $\mathrm{Cay}(S_n, S)$ is called the modified bubble-sort graph of dimension n. We denote these generating transpositions by s_i, where $s_1 = (1, 2), s_2 = (2, 3), \ldots, s_n = (n, 1)$.

Lemma 3. *Suppose the transposition graph $T(S)$ is an n-cycle, and let G denote the automorphism group of $\mathrm{Cay}(S_n, S)$. Then, G_e is the semidirect product $L_e \rtimes D_{2n}$, where G_e is the stabilizer in G of e and L_e is the stabilizer in G of e and each of its neighbors.*

Proof: Let ρ and σ denote the rotation $(1,2,\ldots,n)$ and reflection $(1)(2,n)(3,n-1)\ldots$, respectively, that generate the dihedral group $D_{2n}=\langle\rho,\sigma\rangle\leq S_n$. Consider the conjugation map $\psi: S_n\to \mathrm{Sym}(S_n), x\mapsto x^*$, where $x^*: S_n\to S_n, g\mapsto x^{-1}gx$. Then, ψ is an injective group homomorphism since the center of S_n is trivial for $n\geq 3$ (recall that we assumed that $n\geq 4$ since the $n\leq 3$ cases are easily dealt with). Hence, $D_{2n}^*:=\langle\rho^*,\sigma^*\rangle$, which is a subgroup of the group of inner automorphisms of S_n, is isomorphic to D_{2n}. Every element of D_{2n}^*, when viewed as a permutation of the vertices of the Cayley graph $\mathrm{Cay}(S_n,S)$, fixes the vertex e corresponding to the identity group element in S_n. Thus, the neighbors of the vertex e in the Cayley graph, namely the set of vertices S, is fixed setwise by each element in D_{2n}^*. In fact, ρ^* maps $s_1\mapsto s_2, s_2\mapsto s_3,\ldots,s_n\mapsto s_1$, and σ^* interchanges s_1 and s_n, interchanges s_2 and s_{n-1}, and so on. Thus, $D_{2n}^*\leq \mathrm{Aut}(S_n,S)$, and hence $D_{2n}^*\leq G_e$.

Since $L_e\leq G_e$, we have $D_{2n}^*L_e\leq G_e$. We now show that $G_e\leq D_{2n}^*L_e$. Let $\pi\in G_e$. Since π fixes e, it induces a permutation of its neighbors S. Let τ be the restriction of π to S. We claim that τ is an element of the dihedral group that acts as a group of automorphisms of the n-cycle graph with vertex set $\{s_1,\ldots,s_n\}$. To prove this, suppose τ maps s_1 to s_i. Since s_2 and s_n are the only edges in $T(S)$ incident to s_1, by Lemma 2 there are only two elements s_j of S that do not lie in unique 4-cycles passing through e, s_1 and s_j, namely the elements s_2 and s_n. Similarly, the only two elements s_k of S that do not lie in unique 4-cycles passing through e, s_i and s_k are the elements s_{i-1} and s_{i+1}. Thus, τ maps the set $\{s_2,s_n\}$ to $\{s_{i-1},s_{i+1}\}$. If τ maps s_2 to s_{i+1}, by a similar reasoning, it maps s_3 to s_{i+2} or s_i, but since s_1 is mapped to s_i, s_3 is mapped to s_{i+2}. Continuing in this manner, we see that the action of τ on S is that of a rotation that maps s_i to s_{i+1}. If τ maps s_2 to s_{i-1}, then by a similar reasoning, it maps s_3 to s_{i-2}, s_n to s_{i+1}, and so on, effecting the action of reflection on S. Thus, the action of any element of G_e on S is the same as the action of some element of the dihedral group obtained by restricting D_{2n}^* to S.

For any such τ, since the restriction of D_{2n}^* to S is closed under inverses, there exists an element $\alpha\in D_{2n}$ such that the action of the conjugation map $\alpha^*\in D_{2n}^*$ when restricted to S is the same as the action of τ^{-1} on S. Thus, the action of $\alpha^*\pi\in G_e$ on $\{e\}\cup S$ is the identity, i.e. $\alpha^*\pi\in L_e$. Hence, $\pi\in D_{2n}^*L_e$ and $G_e\leq D_{2n}^*L_e$.

Since $G_e=D_{2n}^*L_e$, $D_{2n}^*\cap L_e=1$, and by Biggs [1, Chapter 16] L_e is a normal subgroup of G_e, we get that $G_e=L_e\rtimes D_{2n}^*$. ∎

Lemma 4. *(Feng [3, Corollary 2.5]) Let S be a set of transpositions generating S_n satisfying the following condition for any two distinct transpositions $t,k\in S$: $tk=kt$ if and only if $\mathrm{Cay}(S_n,S)$ has a unique 4-cycle containing e,t and k, and if $tk\neq kt$ then $\mathrm{Cay}(S_n,S)$ has a unique 6-cycle containing e,t,k and a vertex at distance 3 from e. Then $\mathrm{Cay}(S_n,S)$ is normal and has automorphism group isomorphic to $R(S_n)\rtimes \mathrm{Aut}(S_n,S)$.*

Theorem 5. *Suppose the transposition graph $T(S)$ is the n-cycle graph, and let $n\geq 4$. Suppose $s_i,s_j\in S$ are adjacent transpositions in $T(S)$, and let $\Gamma:=\mathrm{Cay}(S_n,S)$. Then, there exists a unique 6-cycle in Γ containing e,s_i,s_j and some vertex of distance 3 from e if and only if $n\neq 4$.*

Proof: Let $s_i = (i, i+1)$ and $s_{i+1} = (i+1, i+2)$ be two adjacent edges of the n-cycle graph $T(S)$. We take the subscripts to be modulo n; for example, $s_i = (n, 1)$ and $s_{i+1} = (1, 2)$ is possible. The sequence of 6 distinct vertices

$$C := (e, s_i, s_{i+1}s_i, s_is_{i+1}s_i = s_{i+1}s_is_{i+1} = (i, i+2), s_is_{i+1}, s_{i+1}, e)$$

forms a 6-cycle in Γ. Also, the middle vertex $s_is_{i+1}s_i = (i, i+2)$ is not adjacent to e since $T(S)$ does not have triangles, and the distance from $(i, i+2)$ to e is odd because Γ is bipartite and $(i, i+2)$ is an odd permutation. Hence, the distance from $(i, i+2)$ to e is equal to 3. Thus, Γ has at least one 6-cycle containing e, s_i, s_{i+1} and some vertex at distance 3 from e. We now show that the condition that the 6-cycle contain e, s_i, s_{i+1} and a vertex at distance 3 from e forces the uniqueness of this 6-cycle if and only if $n \neq 4$.

Let $(e, s_i, x, w, y, s_{i+1}, e)$ be a 6-cycle in Γ. Then there exist integers k, r, s and ℓ such that $x = s_ks_i$, $w = s_rs_ks_i = s_ss_\ell s_{i+1}$, and $y = s_\ell s_{i+1}$, where w has distance 3 from e. The condition that the six vertices of the 6-cycle be distinct forces $k \neq i, r \neq k, s \neq r, \ell \neq s, \ell \neq i+1$. We determine whether there is any other value of k, r, s and ℓ besides $k = i+1, r = i, \ell = i, s = i+1$ satisfying the two conditions that $s_{i+1}s_is_k = s_\ell s_ss_r$ and the distance from $w = s_rs_ks_i = s_ss_\ell s_{i+1}$ to e is 3.

We consider a few cases, depending on the value of k:

Case 1: Suppose that $k \notin \{i-1, i, i+1, i+2\}$. Since $s_{i+1}s_i = (i, i+1, i+2)$ has disjoint support from $(k, k+1)$, $s_{i+1}s_is_k = s_\ell s_ss_r$ implies that $s_\ell s_ss_r$ is also the product of a transposition $(k, k+1)$ and a three-cycle permutation $(i, i+1, i+2)$. Since $r \neq k$, either $\ell = k$ or $s = k$.

Suppose $\ell = k$. Then $s_ss_r = (i, i+1, i+2)$. If the product of two transpositions is a permutation consisting of a single three-cycle, then the two transpositions must have overlapping support whose union is equal to the three elements of the cycle. Furthermore, since $T(S)$ is the n-cycle graph and $n \geq 4$, $T(S)$ does not contain any triangles, and so $(i, i+2) \notin S$. This forces the equation $s_ss_r = (i, i+1, i+2)$ to have the unique solution $s = i+1, r = i$. But then $w = s_rs_ks_i = s_is_ks_i = s_k$ has distance 1 from e, a contradiction.

Suppose $s = k$. Then, we again have that $s_\ell s_r$ is the three-cycle $(i, i+1, i+2)$. As before, this implies that $\ell = i+1$ and $r = i$, a contradiction because $l \neq i+1$.

Case 2: Suppose $k = i-1$. From the equations $s_{i+1}s_is_k = s_\ell s_ss_r$ and $s_k = s_{i-1}$, we get $(i-1, i, i+1, i+2) = s_\ell s_ss_r$. If the product of three transpositions is a permutation consisting of a single four-cycle, then the transposition graph of these three transpositions is a spanning tree on the same four vertices whose support is the four-cycle. The transposition graph $T(S)$ is the n-cycle graph, and so if $n \geq 5$, $T(S)$ has a unique spanning tree on the vertex set $\{i-1, i, i+1, i+2\}$, namely the path graph containing the three edges $(i-1, i), (i, i+1)$ and $(i+1, i+2)$; however, if $n = 4$, then $(i-1, i+2) \in S$ is also an edge in $T(S)$, which yields other spanning trees in $T(S)$, such as the tree consisting of the three edges $(i-1, i+2), (i+1, i+2)$ and $(i, i+1)$.

If $n \geq 5$, the unique spanning tree given above forces $\{s_{i+1}, s_i, s_k\} = \{s_\ell, s_s, s_r\}$; furthermore, the equation $(i-1, i, i+1, i+2) = s_\ell s_ss_r$ implies that $\ell = i+1$. But this is a contradiction because $\ell \neq i+1$. Thus, if $n \geq 5$, there are no 6-cycles satisfying $k = i-1$.

If $n = 4$, then $(i-1, i+2) \in S$, the permutation $(i-1, i, i+1, i+2)$ can also be expressed as the product $(i-1, i+2)(i+1, i+2)(i, i+1)$; thus the solution $\ell = i+2, s = i+1, r = i$ also satisfies the two conditions $s_{i+1}s_is_k = s_\ell s_s s_r$ and the distance from w to e is 3. Thus, the 6-cycle

$$C' := (e, s_i, s_{i-1}s_i, s_is_{i-1}s_i = s_{i+1}s_{i+2}s_{i+1}, s_{i+2}s_{i+1}, s_{i+1}, e)$$

also satisfies the conditions in the assertion; here $(i-1, i+1) = (i+1, i+3)$ since $i-1$ and $i+3$ are identified with the same vertex of the 4-cycle.

Thus, when $n = 4$, there exists more than one 6-cycle in Γ that contains e, s_i, s_{i+1} and a vertex at distance 3 from e. Since non-uniqueness has now been established for $n = 4$, for the rest of the proof we may assume that $n \geq 5$.

Case 3: Suppose $k = i+2$. Then, $s_{i+1}s_is_k = (i, i+1, i+3, i+2)$. As before, when $n \geq 5$, $T(S) = C_n$ has a unique spanning tree containing any four consecutive vertices. The equation $(i, i+1, i+3, i+2) = s_\ell s_s s_r$ implies $\{s_\ell, s_s, s_r\} = \{s_i, s_{i+1}, s_{i+2}\}$. However, the only way to obtain the product $(i, i+1, i+3, i+2)$ using these three generating transpositions is by having the first term s_ℓ equal s_{i+1}. But this is a contradiction because $\ell \neq i+1$.

Case 4: Suppose $k = i+1$. Here, we have $s_{i+1}s_is_{i+1} = s_\ell s_s s_r$. Again, since $n \geq 5$, the only way to achieve the product $s_{i+1}s_is_{i+1} = (i, i+2)$ using three transpositions from $\{s_i, s_{i+1}, s_{i+2}\}$ is either by the product $s_is_{i+1}s_i$ or $s_{i+1}s_is_{i+1}$. Since $\ell \neq i+1$, we get $\ell = i, s = i+1$ and $r = i$, yielding the unique 6-cycle $C = (e, s_i, s_{i+1}s_i, s_is_{i+1}s_i = s_{i+1}s_is_{i+1} = (i, i+2), s_is_{i+1}, s_{i+1}, e)$ given earlier.

Also, observe that when $k = i+1$ and $n = 4$, we have $(i+3, i) \in S$, and so the product $(i, i+2)$ has other decompositions such as $s_{i+3}s_{i+2}s_{i+3}$. Thus we obtain the solution $\ell = i+3, s = i+2, r = i+3 = i-1$, and another 6-cycle

$$C'' := (e, s_i, s_{i+1}s_i, s_{i-1}s_{i+1}s_i = s_{i+2}s_{i-1}s_{i+1}, s_{i-1}s_{i+1}, s_{i+1}, e).$$

We have shown that: if $n \geq 5$, then $k = i+1$ and there is a unique 6-cycle C in Γ containing e, s_i, s_{i+1} and a vertex at distance 3 from e; if $n = 4$, then there exist other 6-cycles besides C, such as C' and C''. ∎

In the proof above, it was sufficient to just show the existence of three 6-cycles C, C' and C'' since non-uniqueness was all we needed to establish. A consideration of all possible solutions of the equation $s_{i+1}s_is_k = s_\ell s_s s_r$ in the proof above gives the following result:

Lemma 6. *If $s_i, s_{i+1} \in S$ are two adjacent edges in $T(S) = C_4$, then there are exactly eight distinct 6-cycles in $\mathrm{Cay}(S_4, S)$ that contain e, s_i, s_{i+1} and a vertex at distance 3 from e. The total number of vertices in 6-cycles of $\mathrm{Cay}(S_4, S)$ that contain e, s_i and s_{i+1} and that are at distance 3 from e is exactly 6.*

Proof: Let $(e, s_i, s_k s_i, s_r s_k s_i = s_s s_\ell s_{i+1}, s_\ell s_{i+1}, s_{i+1}, e)$ be a 6-cycle containing e, s_i, s_{i+1} and a vertex at distance 3 from e. Since these 6 vertices of the 6-cycle must be distinct, we require that $k \neq i, r \neq k, s \neq r, \ell \neq s, \ell \neq i+1$. As in the proof above, the number of solutions to the equation $s_\ell s_s s_r = s_{i+1} s_i s_k$ is examined for different values of k.

As in the proof of Theorem 5, we consider different cases depending on the value of k:

Case 1: Suppose $k \notin \{i-1, i, i+1, i+2\}$. It has been shown in the proof above that there are no 6-cycles in this case.

Case 2: Suppose $k = i-1$. If the product of three transpositions is a 4-cycle, then these three transpositions form a spanning tree in the transposition graph. Thus, $s_\ell s_s s_r = s_{i+1} s_i s_{i-1} = (i-1, i, i+1, i+2)$ implies that the set of edges $\{s_\ell, s_s s_r\}$ of $T(S)$ forms a spanning tree, and hence a path subgraph of $T(S) = C_4$. Thus, the four possible solutions to the equation $s_\ell s_s s_r = s_{i+1} s_i s_{i-1}$ are $s_\ell = s_{i+1}, s_s = s_i, s_r = s_{i-1}$, and the three cyclic shifts of this solution, i.e. (s_ℓ, s_s, s_r) can also equal $(s_{i+2}, s_{i+1}, s_i), (s_i, s_{i-1}, s_{i+2})$ or $(s_{i-1}, s_{i+2}, s_{i+1})$. The first of these four solutions is not possible since $\ell \neq i+1$. The latter three solutions give rise to the following three 6-cycles:

$\gamma_1 = (e, s_i, s_{i-1} s_i, s_i s_{i-1} s_i = s_{i+1} s_{i+2} s_{i+1}, s_{i+2} s_{i+1}, s_{i+1}, e)$
$\gamma_2 = (e, s_i, s_{i-1} s_i, s_{i+2} s_{i-1} s_i = s_{i-1} s_i s_{i+1}, s_i s_{i+1}, s_{i+1}, e)$
$\gamma_3 = (e, s_i, s_{i-1} s_i, s_{i+1} s_{i-1} s_i = s_{i+2} s_{i-1} s_{i+1}, s_{i-1} s_{i+1}, s_{i+1}, e)$

Case 3: Suppose $k = i+2$. In this case, the only solutions to $s_\ell s_s s_r = s_{i+1} s_i s_{i+2} = (i, i+2, i+3, i+2)$ satisfying $\ell \neq i+1$ are (s_ℓ, s_s, s_r) equals (s_{i+2}, s_i, s_{i+3}) or (s_i, s_{i+2}, s_{i+3}). This gives the two 6-cycles

$\gamma_4 = (e, s_i, s_{i+2} s_i, s_{i+3} s_{i+2} s_i = s_i s_{i+2} s_{i+1}, s_{i+2} s_{i+1}, s_{i+1}, e)$
$\gamma_5 = (e, s_i, s_{i+2} s_i, s_{i+3} s_{i+2} s_i = s_{i+2} s_i s_{i+1}, s_i s_{i+1}, s_{i+1}, e)$

Case 4: Suppose $k = i+1$: We require that $s_\ell s_s s_r = s_{i+1} s_i s_{i+1} = (i, i+2)$. The only way to express $(i, i+2)$ as a product of three transpositions from $\{s_i, s_{i+1}, s_{i+2}, s_{i+3}\}$ is as $s_{i+1} s_i s_{i+1}, s_i s_{i+1} s_i, s_{i+3} s_{i+2} s_{i+3}$ or $s_{i+2} s_{i+3} s_{i+2}$. Since $\ell \neq i+1$, only the latter three products are possible. From these three products, we obtain the following three 6-cycles:

$\gamma_6 = (e, s_i, s_{i+1} s_i, s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, s_i s_{i+1}, s_{i+1}, e)$
$\gamma_7 = (e, s_i, s_{i+1} s_i, s_{i+3} s_{i+1} s_i = s_{i+2} s_{i+3} s_{i+1}, s_{i+3} s_{i+1}, s_{i+1}, e)$
$\gamma_8 = (e, s_i, s_{i+1} s_i, s_{i+2} s_{i+1} s_i = s_{i+3} s_{i+2} s_{i+1}, s_{i+2} s_{i+1}, s_{i+1}, e)$

Thus, there are exactly eight distinct 6-cycles containing e, s_i, s_{i+1} and a vertex at distance 3 from e. The number of vertices in these 8 cycles that are at distance 3 from e is exactly 6 since the middle vertices in γ_4 and γ_5 are the same, and the middle vertices in γ_3 and γ_7 are the same (the vertex $i-1$ is identified with $i+3$ since $T(S) = C_4$, and s_{i+1} and s_{i-1} commute). ∎

Theorem 7. *Let S be a set of transpositions in S_n such that the transposition graph $T(S)$ is the n-cycle graph. Let L_e denote the stabilizer of a vertex neighborhood in* $\mathrm{Cay}(S_n, S)$. *Then, L_e is trivial if and only if $n \neq 4$, and L_e is isomorphic to the Klein 4-group $\mathbb{Z}_2 \times \mathbb{Z}_2$ if $n = 4$.*



Figure 1: The Cayley graph $\mathrm{Cay}(S_4, S)$ when the transposition graph $T(S)$ is a 4-cycle.

Proof: Let $T(S)$ be the n-cycle graph, and let L_e denote the set of automorphisms of $\Gamma := \mathrm{Cay}(S_n, S)$ that fixes the vertex e and each of its neighbors. Define $\Gamma_i(e)$ to be the set of vertices of Γ that have distance i to vertex e.

First consider the case $n = 4$. In this case, $\mathrm{Cay}(S_4, S)$ is the graph shown in Figure 1. The figure of this graph can be obtained by, say, letting vertex 1 represent the identity element e in S_4, and letting vertices $2, 3, 4$ and 5 correspond to the generators s_1, s_2, s_3 and s_4, respectively. For example, vertex 11 corresponds to the element $s_2s_4 = s_4s_2$. We now prove that $L_e \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

Let $\pi \in L_e$. Thus, the action of π on the vertex set $\{1, 2, 3, 4, 5\}$ is the identity. By Lemma 2, there is a unique 4-cycle in Γ containing vertices 1, 4 and 2, namely the cycle $(1, 4, 6, 2, 1)$, and a unique 4-cycle containing 1, 3 and 5, namely the cycle $(1, 3, 11, 5, 1)$. Hence, since $\pi \in L_e$ fixes each s_i, π fixes 6 and 11 also. Since 24 is the unique vertex at maximum distance from 1, π fixes 24 also. Note that π fixes 4 and hence permutes its neighbors $\{1, 12, 6, 13\}$, of which π fixes 1 and 6. Hence, $\{12, 13\}$ is a fixed block of π. Similarly, we get that $\{9, 10\}$, $\{7, 8\}$ and $\{14, 15\}$ are also fixed blocks of π. Also, the vertices in $\Gamma_3(1)$ that are not neighbors of 24, i.e. the vertices $\{16, 17, 21, 22\}$ also form a fixed block of π.

Now π either fixes $\{12, 13\}$ pointwise or interchanges 12 and 13. Thus, π either fixes their neighbors in $\Gamma_3(1)$ (i.e. the sets $\{16, 18, 17\}$ and $\{21, 22, 18\}$ are each fixed blocks of π) or π induces a bijection between these two sets. In either case, 18 is the only vertex from these two sets that is adjacent to 24. Hence, π fixes 18. By a similar argument (that π either fixes $\{9, 10\}$ pointwise or interchanges 9 and 10), we get that π fixes 20. The neighbors of 6 in $\Gamma_3(1)$ form a fixed block of π, and so π fixes $\{19, 20\}$ and hence fixes 19 as well. Since the neighbors of 11 are $\{3, 5, 18, 23\}$ and π fixes 11 and three of its neighbors, π also fixes the remaining neighbor 23. Thus, if $\pi \in L_e$, then the action of π on $\{1, 2, 3, 4, 5, 6, 11, 18, 19, 20, 23\}$ is the identity.

Now $\{12, 13\}$ is a fixed block of π. We consider 2 cases, depending on whether π interchanges 12 and 13 or whether π fixes $\{12, 13\}$ pointwise:

Case 1: Suppose π interchanges 12 and 13. Then π induces a bijection between the neighbors $\{16, 18, 17\}$ of 12 and the neighbors $\{18, 21, 22\}$ of 13. Since π fixes 18, it induces a bijection between $\{16, 17\}$ and $\{21, 22\}$.

Case 1.1: Suppose π maps 16 to 21. Then $\pi : 17 \mapsto 22$. So π maps the neighbors $\{16, 9, 8, 14\}$ of 16 to the neighbors $\{13, 9, 7, 14\}$ of 21. But $\{9, 10\}$ is a fixed block of π, hence $\pi : 9 \mapsto 9, 8 \mapsto 7, 7 \mapsto 8, 14 \mapsto 14, 15 \mapsto 15$. Since the neighbors $\{16, 21, 20\}$ of 9 are permuted by π, we have $\pi : 21 \mapsto 16$. Hence $\pi : 22 \mapsto 17$. Thus, $\pi = (12, 13)(16, 21)(22, 17)(7, 8)$. Denote this permutation by $\pi_1 := (12, 13)(16, 21)(22, 17)(7, 8)$.

Case 1.2: Suppose $\pi : 16 \mapsto 22$. Hence $\pi : 17 \mapsto 21$. So π maps the neighbors $\{12, 9, 8, 14\}$ of 16 to the neighbors $\{13, 10, 7, 15\}$ of 22. By the conditions above, $\pi : 12 \mapsto 13, 9 \mapsto 10$ (hence $10 \mapsto 9$), $8 \mapsto 7$ (hence $7 \mapsto 8$), and $14 \mapsto 15$ (hence $15 \mapsto 14$). Since π interchanges 7 and 8, π induces a bijection between their neighbors $\{21, 22, 23\}$ and $\{16, 17, 23\}$. Since π fixes 23, it induces a bijection between $\{21, 22\}$ and $\{16, 17\}$. We know $\pi : 16 \mapsto 22, 17 \mapsto 21$. We consider two subcases:

Case 1.2.1: $\pi : 22 \mapsto 17, 21 \mapsto 16$. So the permutation π contains $(16, 22, 17, 21)$ as a cycle. In this case, π maps the neighbors $\{12, 9, 8, 14\}$ of 16 to the neighbors $\{13, 10, 7, 15\}$ of 22 and the neighbors of 22 to the neighbors $\{12, 10, 8, 15\}$ of 17. The first restriction implies $\pi : 9 \mapsto 10$, and the second implies $\pi : 10 \mapsto 10$, an impossibility. Hence, this subcase is not possible.

Case 1.2.2: π interchanges 16 and 22, and it interchanges 17 and 21. In this case, $\pi = (12, 13)(9, 10)(8, 7)(14, 15)(21, 17)(16, 22)$. Denote this permutation by $\pi_2 := (12, 13)(9, 10)(8, 7)(14, 15)(21, 17)(16, 22)$.

Case 2: Suppose $\pi : 12 \mapsto 12, 13 \mapsto 13$. Then the neighbors $\{16, 17\}$ of 12 are a fixed block of π.

Case 2.1: If π fixes 16, then the set of its neighbors $\{12, 9, 8, 14\}$ is a fixed block of π, and hence is fixed by π pointwise since each of 9, 8 and 14 are already known to lie in different fixed blocks of π. Hence, π also fixes 10, 7, and 15. Also, π fixes 17 since it fixes the neighbors $\{16, 18, 17\}$ of 12 and since 16 and 18 are fixed points. Also, the neighbors $\{17, 19, 22\}$ of 15, and hence 22, and the neighbors $\{21, 19, 16\}$ of 14, and hence also 21, are all fixed by π. Thus, π is the identity element.

Case 2.2: If $\pi : 16 \mapsto 17$, then $17 \mapsto 16$. So π induces a bijection between their neighbors $\{12, 9, 8, 14\}$ and $\{12, 10, 8, 15\}$. We know $12 \mapsto 12$, so $9 \mapsto 10$, and hence $10 \mapsto 9$. Similarly, $8 \mapsto 8$ (hence $7 \mapsto 7$), and $14 \mapsto 15, 15 \mapsto 14$. Since $\{16, 17, 21, 22\}$ is a fixed block, $\{21, 22\}$ is also a fixed block. If $\pi : 21 \mapsto 21$, then its neighbors $\{13, 9, 7, 14\}$ form a fixed block, meaning 9 is a fixed point, a contradiction. Hence, $\pi : 21 \mapsto 22, 22 \mapsto 21$, and we get that $\pi = (9, 10)(16, 17)(14, 15)(21, 22)$, which is seen to equal $\pi_1\pi_2$.

Thus, the only elements in L_e are π_1 (Case 1.1), π_2 (Case 1.2.2), the identity element (Case 2.1), and $\pi_1\pi_2$ (Case 2.2). The order of π_1 and π_2 is 2; thus, L_e is isomorphic to the Klein 4-group $\mathbb{Z}_2 \times \mathbb{Z}_2$.

Now consider the case $n \geq 5$. Let $\pi \in L_e$, and let $t, k \in S$. If t and k have disjoint support, then by Lemma 2 there is a unique 4-cycle in Γ containing e, t and k. Since π is an automorphism of Γ that fixes e, t and k, and since tk is the only other common neighbor of t and k besides e, π fixes tk also. If t and k have overlapping support, then by Theorem 5 there is a unique 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e. Hence, any automorphism of Γ fixing e, t and k must also fix this 6-cycle (and hence fix this 6-cycle pointwise since it fixes three of the elements in this 6-cycle), which contains tk. Thus, π fixes tk. Thus, if $\pi \in L_e$ and $s_1, s_2 \in S$, then π also fixes $s_1 s_2$.

It now suffices to prove that π fixes an arbitrary vertex $s_1 s_2 \cdots s_k$, and this can be shown by induction. Assume this assertion holds for smaller values of k. Define $z = s_3 \cdots s_k$, and let $r_z : S_n \to S_n, g \mapsto gz$ be right translation by z. Then $\psi := r_z \pi r_{z^{-1}}$ is a composition of automorphisms of the Cayley graph Γ and hence an automorphism of Γ. By the inductive hypothesis, $e^\psi = e^{r_z \pi r_{z^{-1}}} = z^{\pi r_{z^{-1}}} = z^{r_{z^{-1}}}$, which equals e. Hence, ψ fixes e. Also, ψ fixes $s_i \in S$ because $s_i^\psi = s_i^{r_z \pi r_{z^{-1}}} = (s_i s_3 \cdots s_k)^{\pi r_{z^{-1}}} = (s_i s_3 \cdots s_k)^{r_{z^{-1}}}$ (by the inductive hypothesis), which equals s_i. Thus, $\psi \in L_e$, and by the previous paragraph $(s_1 s_2)^\psi = s_1 s_2$. But this is equivalent to the assertion that ψ fixes the element $s_1 \ldots s_k$ because $(s_1 s_2)^{\rho_z \pi \rho_{z^{-1}}} = s_1 s_2$ iff $(s_1 s_2 z)^\pi = (s_1 s_2 z)$. ∎

By the results proved thus far, it follows that:

Corollary 8. *Let S be a set of transpositions in S_n. If the transposition graph $T(S)$ is a 4-cycle, then the automorphism group of the Cayley graph* $\mathrm{Cay}(S_4, S)$ *is isomorphic to $R(S_4)G_e = R(S_4)(L_e \rtimes D_8) = R(S_4)(V_4 \rtimes D_8)$, and $R(S_4)$ is not a normal subgroup of the automorphism group of* $\mathrm{Cay}(S_4, S)$. *If $T(S)$ is the n-cycle graph, where $n \geq 5$, then the automorphism group of* $\mathrm{Cay}(S_n, S)$ *is the semidirect product $R(S_n) \rtimes D_{2n}$.*

Theorem 9. *Let S be a set of transpositions generating S_n, and let Γ be the Cayley graph* $\mathrm{Cay}(S_n, S)$. *Let $k, t \in S$ be distinct transpositions. If $tk \neq kt$ and the girth of the transposition graph $T(S)$ is at least 5, then there is a unique 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e. Moreover, if $T(S)$ does contain a 4-cycle, then there exist $t, k \in S$ such that $tk \neq kt$ and such that there does not exist a unique 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e.*

Proof: Suppose $T(S)$ does not contain triangles or 4-cycles, and let $t, k \in S$ be distinct transpositions such that $tk \neq kt$. We show that there is a unique 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e. Since $tk \neq kt$, t and k are adjacent edges in $T(S)$, so we may assume without loss of generality that $k = (1, 2)$ and $t = (2, 3)$. Then, $(e, t, kt, tkt = ktk, tk, k, e)$ is a 6-cycle in Γ. The distance from e to tkt is at most 3. Also, Γ is a bipartite graph with bipartition equal to the even and odd permutations, and so the distance from e to tkt is odd. But this distance cannot be 1 since $tkt = (1, 3) \notin S$ because $T(S)$ has no triangles. Thus, tkt has distance 3 to e. We show next that this 6-cycle is the unique 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e.

Suppose $(e, t, t_1t, t_2t_1t = k_2k_1k, k_1k, k, e)$ is a 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e. Since the 6 vertices in this 6-cycle are distinct, we have that $t_1 \neq t, t_2 \neq t_1, k_2 \neq t_2, k_2 \neq k_1$ and $k_1 \neq k$. We first show that $k_1 = t = (2, 3)$. Suppose $k_1 \neq (2, 3)$. We know $k_1 \neq k = (1, 2)$ and since $T(S)$ has no triangles, $k_1 \neq (1, 3)$. Thus, $\{1, 2, 3\}$ and the support of k_1 have an intersection of size at most 1. (For example, if $k_1 = (1, 4)$ say, then the intersection is $\{1\}$, and if $k_1 = (4, 5)$ say, then the intersection is trivial.) This implies that the union of $\{1, 2, 3\}$ and the support of k_1 is at least 4 because k_1 contributes at least one new element to this union. By $t_2t_1t = k_2k_1k$, we have $k_2t_2t_1 = k_1kt$. We now prove that the assumption that $T(S)$ have girth at least 5 forces $\{k_2, t_2, t_1\} = \{k_1, k, t\}$. Now $k_1 \neq (2, 3)$ since $T(S)$ does not contain triangles, and k and t have total support $\{1, 2, 3\}$.

Suppose k_1 has overlapping support with $\{1, 2, 3\}$. This can happen if and only if the three edges $\{k_1, k, t\}$ form a tree spanning 4 distinct vertices of $T(S)$, namely the vertices $\{1, 2, 3, i\}$ for some i. The product k_1kt of three transpositions that form a subtree in $T(S)$ is a permutation that consists of a single four-cycle, and conversely (cf. Godsil and Royle [5, Lemma 3.10.2]), if the product of three transpositions is a four-cycle, then these 3 transpositions constitute the edges of a spanning tree on the same 4 vertices corresponding to the support of the four-cycle. Thus, the edges $\{k_2, t_2, t_1\}$ also form a spanning tree on the same 4 vertices $\{1, 2, 3, i\}$. Now, the only spanning tree subgraph in $T(S)$ on these 4 vertices is the spanning tree $\{k_1, k, t\}$ because $T(S)$ does not contain triangles or 4-cycles. Thus, $\{k_2, t_2, t_1\} = \{k_1, k, t\}$. The rest of this proof of the uniqueness of 6-cycles follows the proof in [3, p. 71].

If k_1 has disjoint support from $\{1, 2, 3\}$, then the product k_1kt has a 2-cycle, say (i, j) and a 3-cycle $kt = (1, 3, 2)$, the two cycles being disjoint. By $k_2t_2t_1 = k_1kt$, we have that at least one of k_2, t_2 or t_1 must equal (i, j), and the product of the other two must equal $(1, 3, 2)$. Since $T(S)$ has no triangles, the only way to decompose $(1, 3, 2)$ as a product of two transpositions from S is as $(1, 2)(2, 3) = kt$. Thus, two of k_2, t_2, t_1 must equal k and t, and so we again get $\{k_2, t_2, t_1\} = \{k_1, k, t\}$.

Thus, if $T(S)$ does not contain triangles or 4-cycles and $k_1 \neq (2, 3)$, then the union of the support of k_1, k and t is at least 4, and hence $k_2t_2t_1 = k_1kt$ forces $\{k_2, t_2, t_1\} = \{k_1, k, t\}$. A few cases now arise, each of which is seen to be impossible:

We already have $k_2 \neq k_1$. We consider the two cases $k_2 = k$ and $k_2 = t$:

Case 1: $k_2 = k$. Then, either $t_2 = k_1$ and $t_1 = t$ or $t_2 = t$ and $t_1 = k_1$. The first subcase is impossible because $t_1 \neq t$. The second subcase $t_2 = t$ and $t_1 = k_1$, along with $k_2t_2t_1 = k_1kt$ gives $ktk_1 = k_1kt$, i.e. $k_1 = (1, 2, 3)k_1(1, 3, 2)$. This equation implies that the support of k_1 is disjoint from $\{1, 2, 3\}$, and so $k_2k_1k = k_2kk_1 = k_1 \in S$, implying that the vertex k_2k_1k is at distance 1 and not distance 3 from e, a contradiction.

Case 2: $k_2 = t$: Then we have two subcases - that either $t_2 = k_1$ and $t_1 = k$ or $t_2 = k$ and $t_1 = k_1$. In the first subcase, $k_2t_2t_1 = k_1kt$ gives $k_1 = (2, 3)k_1(1, 3)$. This equation has no solutions. For if 1 is not in the support of k_1, then k_1 doesn't move 1 but the right side does. Since k_1 moves 1, 2 and 3 are not in the support of k_1 (because $k_1 \neq (1, 2), (1, 3)$). So $k_1 = (2, 3)k_1(1, 3) = k_1(2, 3)(1, 3) = k_1(1, 3, 2)$, an impossibility. In the second subcase, we get $k_1 = (1,3,2)k_1(1,3,2)$. This equation also has no solutions. For if the support of k_1 is disjoint from $\{1,2,3\}$, then the right side becomes $k_1(1,2,3)$, which cannot equal k_1. And if the support of k_1 overlaps with $\{1,2,3\}$, say $k_1 = (1,i)$, then the left side maps i to 1, whereas the right side maps i to 3.

Thus, $k_1 = (2,3) = t$.

In a similar manner, it can be shown that $t_1 = (1,2) = k$. For if $t_1 \neq (1,2)$, then since $t_1 \neq t = (2,3)$ and since $t_1 \neq (1,3)$ ($T(S)$ has no triangles), we have that the union of the support of t_1, k and t is at least 4. As before, if $T(S)$ does not have triangles or 4-cycles, then the equation $t_2k_2k_1 = t_1tk$ implies $\{t_2, k_2, k_1\} = \{t_1, t, k\}$. We already have $t_2 \neq t_1$. We consider the two cases $t_2 = t$ and $t_2 = k$ separately:

Case 1: $t_2 = t$: The subcase $k_2 = t_1, k_1 = k$ is not possible since $k_1 \neq k$. If $k_2 = k$ and $k_1 = t_1$, then $k_2k_1k = t_2t_1t$ gives $t_1 = (1,3,2)t_1(1,2,3)$. Thus, t_1 has support disjoint from $\{1,2,3\}$, giving that $t_2t_1t = t_2tt_1 = ttt_1 = t_1 \in S$, a contradiction because the vertex t_2t_1t is required to be at distance 3 from e.

Case 2: $t_2 = k$. Consider first the subcase $k_2 = t_1, k_1 = t$. By $k_2k_1k = t_2t_1t$, we get $t_1tk = kt_1t$, i.e. $t_1 = kt_1tkt = (1,2)t_1(1,3)$. This equation has no solutions, for if the support of t_1 is disjoint from $\{1,2,3\}$, then the equation becomes $t_1 = t_1(1,2)(1,3)$, which is impossible. If t_1 has support overlapping with $\{1,2,3\}$, say $t_1 = (1,i)$ ($i \neq 1,2,3$), then the left side and right side map i to different points. Similarly, $t_1 \neq (2,i),(3,i)$. The second subcase is $k_2 = t, k_1 = t_1$. Then, by $k_2k_1k = t_2t_1t$, we get $tt_1k = kt_1t$, or $t_1 = (1,2,3)t_1(1,2,3)$. Again, in each of the cases where t_1 has support overlapping with $\{1,2,3\}$ and support disjoint from $\{1,2,3\}$, it is seen that the equation has no solutions.

This proves that $t_2 = (1,2)$. Thus, $k_2k_1k = t_2t_1t$ becomes $k_2tk = t_2kt$, or $t_2k_2 = ktkt = (1,2,3)$. Since $T(S)$ has no triangles, the only way to express $(1,2,3) = tk$ as a product of two transpositions t_2k_2 is as $t_2 = t, k_2 = k$. This gives rise to the same 6-cycle given initially, and hence there exists only 6-cycle in Γ containing e, t, k and a vertex at distance 3 from e.

To prove the second part of the theorem, note that if $T(S)$ contains a 4-cycle, then $\Gamma = \mathrm{Cay}(S_n, S)$ contains the modified bubble-sort graph MBS_4 as a subgraph; here MBS_4 denotes the Cayley graph of the group S_4 generated by 4 cyclically adjacent transpositions. If t and k are adjacent transpositions in the 4-cycle $T(S)$, then by Theorem 5, the subgraph MBS_4 has more than one 6-cycle containing e, t, k and a vertex at distance 3 from e. Thus, the Cayley graph $\mathrm{Cay}(S_n, S)$, which contains MBS_4 as a subgraph, also has more than one 6-cycle containing e, t, k and a vertex at distance 3 from e. ∎

From Lemma 4 and Theorem 9, it follows that a sufficient condition for normality of $\mathrm{Cay}(S_n, S)$ is that the transposition graph $T(S)$ have girth at least 5:

Corollary 10. *Let S be a set of transpositions generating S_n such that the girth of the transposition graph $T(S)$ is at least 5. Then, the automorphism group of the Cayley graph* $\mathrm{Cay}(S_n, S)$ *is the semidirect product* $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$.

A special case of this Corollary whereby $T(S)$ is a tree gives the result of Feng [3] that the automorphism group of a Cayley graph generated by a transposition tree is isomorphic to $R(S_n) \rtimes \mathrm{Aut}(S_n, S)$.

References

[1] N. L. Biggs. *Algebraic Graph Theory, 2nd Edition.* Cambridge University Press, Cambridge, 1993.

[2] J. D. Dixon and B. Mortimer. *Permutation Groups.* Graduate Texts in Mathematics vol. 163, Springer, 1993.

[3] Y-Q. Feng. Automorphism groups of Cayley graphs on symmetric groups with generating transposition sets. *Journal of Combinatorial Theory Series B*, 96:67–72, 2006.

[4] C. Godsil. On the full automorphism group of the graph. *Combinatorica*, 1:243–256, 1981.

[5] C. Godsil and G. Royle. *Algebraic Graph Theory.* Graduate Texts in Mathematics vol. 207, Springer, New York, 2001.

[6] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.4.12,* `http://www.gap-system.org`. 2008.

[7] Q. Huang and Z. Zhang. On the Cayley graphs of Sym(n) with respect to transposition connections. *submitted to Europ. J. Combin.*

[8] M. Y. Xu. Automorphism groups and isomorphisms of Cayley digraphs. *Discrete Mathematics*, 182:309–319, 1998.

[9] Z. Zhang and Q. Huang. Automorphism groups of bubble sort graphs and modified bubble sort graphs. *Advances in Mathematics (China)*, 34(4):441–447, 2005.